

Light cluster production in $E/A = 61$ MeV $^{36}\text{Ar} + ^{112,124}\text{Sn}$ reactions

R. Ghetti¹, J. Helgesson

School of Technology and Society, Malmö University, S-205 06 Malmö, Sweden

V. Avdeichikov, B. Jakobsson

Department of Physics, Lund University, Box 118, S-22100 Lund, Sweden

N. Colonna, G. Tagliente

INFN and Dip. di Fisica, V. Amendola 173, I-70126 Bari, Italy

H.W. Wilschut, V.L. Kravchuk²

Kernfysisch Versneller Instituut, Zernikelaan 25 NL 9747 AA Groningen, The Netherlands

Experimental kinetic energy distributions and small-angle two-particle correlation functions involving deuterons and tritons are compared for $^{36}\text{Ar} + ^{112,124}\text{Sn}$ collisions at $E/A = 61$ MeV (i.e. for systems similar in size, but with different isospin content). A larger triton yield is observed from the more neutron-rich system, as predicted by IBUU simulations, while the emission times of the light clusters are found to be the same for the two Sn-target systems. For both systems, the time sequence $\tau_d < \tau_p < \tau_t$, is deduced for charged particles emitted from the intermediate velocity source.

PACS number(s): 25.70.Pq

¹Corresponding Author: Roberta Ghetti, School of Technology and Society, Malmö University, Sweden *E-mail address:* roberta.ghetti@ts.mah.se

²Present address: LNL, Legnaro, Italy.

I. INTRODUCTION

Among the challenges in nuclear physics today, is to understand the nuclear interaction for systems with an exotic composition of neutrons and protons (which may occur in the r -process of nucleosynthesis and in neutron stars). One of the topics in this challenge is to understand the isospin dependence of the nuclear equation of state (EOS), and in particular its density dependence [1–6]. In this difficult task many observables are needed to put as many constraints as possible on the nuclear interaction. Among several suggested observables, [7–13], the two-nucleon correlation function has been considered in Refs. [14,15].

Experimentally, isospin effects in two-particle correlation functions were investigated by our group in Ref. [16]. The experimental data were from $E/A = 61$ MeV ^{36}Ar -induced semi-peripheral collisions on isotope-separated $^{112,124}\text{Sn}$ targets. Stronger np correlations were found for the $\text{Ar} + ^{124}\text{Sn}$ system, pointing to a shorter neutron emission time from the more neutron-rich system. Smaller isospin effects were seen also in pp , nd and nt correlation functions [16]. In addition, a study of the emission time sequence of neutrons, protons, and deuterons from the same reactions was carried out by our group in Ref. [17], utilizing the method of the velocity-gated correlation functions. That investigation revealed a sensitiv-

ity of the particle emission time sequence to the isospin content of the emitting source, particularly for intermediate velocity source emission and emission from the target residues.

In this paper we present complementary information to that of Refs. [16,17], from the same experimental data set, focusing on deuterons and tritons. Indeed, also light cluster production is expected to be a sensitive probe for the isospin dependence of the EOS. In the theoretical investigation of Ref. [18] (where IBUU was coupled with a coalescence model for cluster production), it was found that the nuclear symmetry energy significantly affects the production of light clusters in heavy ion collisions. More deuterons and tritons are produced with a stiff nuclear symmetry energy than with a soft. This is because the stiff symmetry energy induces a stronger pressure in the reaction system, and thus causes an earlier emission of neutrons and protons leading to stronger correlations among nucleons, and, consequently, larger yields of deuterons and tritons.

II. EXPERIMENTAL DETAILS

The experiment was performed at KVI (Groningen). The interferometer consisted of 16 CsI(Tl) detectors for light charged particles, mounted in the angular range $30^\circ \leq \theta \leq 114^\circ$, and 32 liquid scintillator neutron detectors, mounted behind the “holes” of the CsI array, in matching positions to provide the np interferometer [19]. Energy thresholds for protons, deuterons, and tritons in the CsI(Tl) detectors were 8, 11, and 14 MeV respectively. Details about the experimental setup and the particle energy determination are given in Refs. [19–21].

From our previous analyses [16,17], including the source analysis of Ref. [21], we have confirmed that the emission of light particles from 61 A MeV (semi-peripheral) heavy ion reactions originates from (at least) three sources: projectile and target residue evaporative sources (statistical evaporation) and intermediate velocity source. The intermediate source represents dynamical emission, which is described either by early nucleon-nucleon collisions or by other pre-equilibrium processes [22–30]. Within the multi-source reaction mechanism described above, composite particles, like deuterons and tri-

tons, are believed to be predominantly emitted from the dynamical emission source [31,32], where they are formed by a coalescence mechanism [21,33].

III. RESULTS

Figure 1 compares the kinetic energy distributions of deuterons and tritons measured in $^{36}\text{Ar}+^{112}\text{Sn}$ and $^{36}\text{Ar}+^{124}\text{Sn}$ collisions. One can notice that the yields of the deuteron spectra are the same for the two Sn-targets, while the triton yield from the more neutron-rich $^{36}\text{Ar}+^{124}\text{Sn}$ system is enhanced (by a factor 1.1 to 1.5) over the whole angular and energy range considered.




FIG. 1. Comparison of the d and t kinetic energy yields measured in $^{36}\text{Ar}+^{124}\text{Sn}$ (dashed lines) and $^{36}\text{Ar}+^{112}\text{Sn}$ (full lines) at the angles indicated in the figure. The yields are arbitrarily shifted in the y-axis as indicated in the figure.

To further investigate the deuteron and triton production, we present the two-particle correlation functions involving deuterons and tritons. The correlation function, $C(\vec{q}, \vec{P}_{tot}) = kN_c(\vec{q}, \vec{P}_{tot}) / N_{nc}(\vec{q}, \vec{P}_{tot})$, is generated by dividing the coincidence yield (N_c) by the yield of non-correlated events (N_{nc}) constructed from the product of the singles distributions [34]. The relative momentum is, $\vec{q} = \mu(\vec{p}_1/m_1 - \vec{p}_2/m_2)$, where μ is the reduced mass, and the total momentum is, $\vec{P}_{tot} = \vec{p}_1 + \vec{p}_2$. The correlation function is normalized to unity at large values of q , ($200 < q < 300$ MeV/c for all particle pairs) where no correlations are expected.

The pp , np , and pd correlation functions have already been presented in Refs. [16,17]. In those analyses, particles emitted by the intermediate velocity source were enhanced by selecting high- and intermediate- P_{tot} pairs, while particles emitted by the target residues were enhanced by selecting low- P_{tot} pairs in the target frame. Further selection of different sources was obtained by ap-

plying angular gates. The pp and np correlation functions were found to be very sensitive to the applied gates, indicating that both neutrons and protons are emitted from several sources. In contrast, deuterons and tritons mainly originate from common sources with a small spread in source velocity ($v_{source} \approx 0.18$ c) [21]. In agreement with those findings, we find, in the present work, that the dd , tt , and dt correlation functions are quite insensitive to applying total-momentum and angular gates. We have therefore chosen to present angle and total-momentum-integrated dd , tt and dt correlation functions in this paper.

Figure 2(a,b) compares the dd (a) and tt (b) correlation functions measured from the two Sn-target systems. One can see that the Coulomb hole dominates both correlations at low values of the relative momentum. For both dd and tt , the width of the Coulomb hole is the same for the two Sn-targets. This feature indicates that the average emission time of deuterons and tritons is the same for the two systems. The width of the Coulomb hole is larger for dd than for tt , which is consistent with a larger average emission time for tritons than for deuterons.




FIG. 2. From $^{36}\text{Ar}+^{124}\text{Sn}$ (open circles) and $^{36}\text{Ar}+^{112}\text{Sn}$ (filled circles), dd (a), tt (b), and dt (d) angle- and total-momentum-integrated correlation functions. A high-total-momentum gate is applied to the pt (c) correlation function.

The dt correlation function in Fig. 2d is very similar for the two Sn-targets. It is governed by strong resonances at $q \approx 10.8$ MeV/c and $q \approx 83.5$ MeV/c, due to the excited states, $E^* = 16.75$ MeV ($\Gamma_{cm} = 0.076$ MeV) and $E^* = 19.8$ MeV ($\Gamma_{cm} = 2.5$ MeV) of ^5He [35].

We now turn to the correlation functions involving also protons. The pd correlation function, that was presented in Refs. [16,17], is characterized by a pronounced anti-correlation at small relative momenta, due to fi-

nal state Coulomb interaction. The anti-correlation is more pronounced for particle pairs selected by high- and intermediate-total-momentum gates, than for pairs selected by low-total-momentum gate.

The pt correlation function, shown in Fig. 2c, contains contributions of excited states of ${}^4\text{He}$ at $q \approx 23.6$ MeV/c ($E^* = 20.21$ MeV, $\Gamma_{cm} = 0.5$ MeV), $q \approx 41.0$ MeV/c ($E^* = 21.01$ MeV, $\Gamma_{cm} = 0.84$ MeV), and $q \approx 53.4$ MeV/c ($E^* = 21.84$ MeV, $\Gamma_{cm} = 2.01$ MeV) [36], while higher excited states of ${}^4\text{He}$ are much broader and not visible in Fig. 2c. The positions of the 1st and 3rd resonances are marked by arrows.

In contrast with dd , tt , and dt , the pd and pt correlation functions are sensitive to applying angle- and total-momentum gates. This is because protons are emitted from several sources [21]. In order to consistently investigate emission from the intermediate velocity source, we have therefore applied a high-total-momentum gate to the pt correlation function (shown in Fig. 2c). This selection generates a correlation function characterized by stronger resonance peaks as compared to those obtained by lower total momentum gates. Inspection of Fig. 2c reveals also a small difference in the correlation functions for the two Sn isotopes. The more neutron-rich system leads to a slightly stronger resonance peak for the higher excited states of ${}^4\text{He}$, while the opposite behavior is seen for the lowest excited state.

Further information about the emission times and chronologies can be obtained by means of particle-velocity-gated correlation functions of non-identical particles [37–39]. This adds an important and novel piece of information to the picture of the reaction mechanism emerging from previous studies [16,17]. For non-identical particles, a and b , we construct the correlation functions $C_a(q)$, gated on pairs with $v_a > v_b$, and $C_b(q)$, gated on pairs with $v_b > v_a$. The particle velocities are calculated in the frame of the emitting source, and the same normalization constant, calculated from the ungated correlation function, is utilized for both C_a and C_b [38]. If two particles a and b are emitted independently from a source, with a later (earlier) than b , then the ratio C_a/C_b will show a peak (dip) in the region of q where there is a correlation, and a dip (peak) in the region of q where there is an anti-correlation.

To correctly interpret the experimental results, it is important to realize that the resonance peaks may a priori have two different origins [40,41]:

1. From processes where an unstable fragment formed in the reaction decays into two particles which are measured in coincidence (e.g. ${}^5\text{He} \rightarrow d + t$). The energy and momentum conservation in the decay determines the location of the resonance peak.
2. From final state interactions between particles emitted independently from a source (e.g. intermediate velocity source, target and projectile residues).

If the two particles are emitted independently, the velocity-gated correlation functions contain information on the time sequence, but this is not the case if the two particles originate from the two-body decay of an unstable fragment emitted in the reaction. In the latter case, the two particle velocities are determined by momentum conservation, and in the rest system of the decaying fragment, the lightest particle will always get the highest velocity. In this case, the velocity-gated correlation function obtained with the condition that the lightest particle has the largest velocity, should exhibit the strongest correlation or anti-correlation (see Ref. [42] for further details). When we in some cases observe the opposite behavior (namely that the gate, where the *heaviest* particle has the largest velocity, leads to a stronger correlation or anti-correlation) we can reliably conclude that this behavior is dominated by a mechanism other than two-body decay. We attribute this effect to the interaction of independently emitted particles, and in this case, we use the velocity-gated correlation function to obtain information on the time sequence of the independently emitted particles.




FIG. 3. From $E/A = 61$ MeV $^{36}\text{Ar} + ^{112}\text{Sn}$ (left column) and ^{124}Sn (right column) collisions, particle-velocity-gated (filled and open circles) pt (a,c) and dt (e,g) correlation functions, and their ratios (b,d,f,h).

The study of the particle-velocity-gated pd correlation function can be found in Ref. [17]. The extracted emission time sequence is that deuterons are, on average, emitted earlier than protons for both Sn-target systems (consistent with the findings of previous experimental investigations [38,39]). Note that this result refers to the average emission time, and it does not exclude that prompt protons are emitted before any deuteron is emitted.

The particle-velocity-gated pt correlation function is shown in Fig. 3a-d. In the momentum region below $q \approx 50$ MeV/c, the enhancement of $C_p(v_p > v_t)$ over $C_t(v_t > v_p)$, expected from momentum conservation of the two-body decay of ^4He , is not observed. This indicates that the behavior of the pt correlation function in this region is dominated by independently emitted pairs. The emission chronology that can be deduced from this region is that protons are emitted earlier than tritons. Inspection of the C_p/C_t ratios (Fig. 3b,d) reveals that the deduced pt emission chronology is qualitatively similar for the two Sn-target systems.

The particle-velocity-gated dt correlation function in Fig. 3e-h, shows an enhancement of the C_d correlation function that is expected from the two-body decays of ^5He . The behavior of the C_d/C_t ratio (Fig. 3f,h) is therefore of delicate interpretation, since it may be dominated by the two-body resonance decays of ^5He , and it does not lead itself to deduce the emission chronology in this case.

IV. SUMMARY

Kinetic energy spectra and two-particle correlation functions of deuterons and tritons have been investigated in $E/A = 61$ MeV $^{36}\text{Ar} + ^{112,124}\text{Sn}$ reactions. The yield of tritons is larger from the more neutron-rich system, while no differences are found in the deuteron yield. The dd , tt and dt correlation functions are rather insensitive to applying angle- and total-momentum gates, supporting the interpretation [21] that the origin of emission of deuterons and tritons is a coalescence mechanism taking place in the intermediate velocity source created in the early dynamical phase. For pd and pt pairs, instead, the emission from the intermediate velocity source can be enhanced by selecting high-total-momentum pairs.

From Ref. [17] and the analysis of the velocity-gated pt correlation function presented here, we find that, for the intermediate velocity source, deuterons are on average emitted before protons, which in turn are emitted before tritons, $\tau_d < \tau_p < \tau_t$. This is also consistent with the behavior of the dd and tt correlation functions.

The isospin effects in deuteron and triton emission, appear to be weak for the studied systems. Apart for

a larger triton yield from the more neutron-rich system, the emission chronology appears to be the same, as well as the average deuteron and triton emission times. One possible interpretation is that the larger yield of tritons in the $^{36}\text{Ar} + ^{124}\text{Sn}$ system is formed in the same amount of time as in the $^{36}\text{Ar} + ^{112}\text{Sn}$ system, because of a larger abundance of neutrons emitted early in the $^{36}\text{Ar} + ^{124}\text{Sn}$ reaction.

Acknowledgements

The author RG wishes to thank the School of Technology and Society of Malmö University.

-
- [1] *Isospin Physics in Heavy-Ion Collisions at Intermediate Energies*, edited by Bao-An Li and W.U. Schröder (Nova Science Publishers, Inc., New York, 2001).
 - [2] H. Müller and B. Serot, Phys. Rev. C **52**, 2072 (1995).
 - [3] M. Colonna, *et al.*, Phys. Rev. C **57**, 1410 (1998).
 - [4] Bao-An Li, Phys. Rev. Lett. **85**, 4221 (2000).
 - [5] Y.G. Ma, *et al.*, Phys. Rev. C **60**, 024607 (1999).
 - [6] J.-Y. Liu *et al.*, Phys. Rev. C **63**, 054612 (2001).
 - [7] Bao-An Li, *et al.*, Phys. Rev. Lett. **78**, 1644 (1997).
 - [8] V. Baran, *et al.*, Nucl. Phys. **A703**, 603 (2002).
 - [9] Bao-An Li and C.M. Ko, Nucl. Phys. **A618**, 498 (1997).
 - [10] H.S. Xu, *et al.* Phys. Rev. Lett. **85**, 716 (2000).
 - [11] W.P. Tan, *et al.*, Phys. Rev. C **64**, 051901(R) (2001).
 - [12] M.B. Tsang, *et al.*, Phys. Rev. Lett. **86**, 5023 (2001).
 - [13] Bao-An Li, *et al.*, Phys. Rev. C **64**, 054604 (2001).
 - [14] L.W. Chen, *et al.*, Phys. Rev. Lett. **90**, 162701 (2003).
 - [15] L.W. Chen, *et al.*, Phys. Rev. C **68**, 014605 (2003).
 - [16] R. Ghetti, *et al.*, Phys. Rev. C **69**, 031605(R) (2004).
 - [17] R. Ghetti, *et al.*, Phys. Rev. C **70**, 034601 (2004).
 - [18] L.W. Chen, *et al.*, Phys. Rev. C **68**, 017601 (2003).
 - [19] R. Ghetti, *et al.*, Nucl. Inst. Meth. **A 516**, 492 (2004).
 - [20] R. Ghetti, *et al.*, Phys. Rev. Lett. **91**, 092701 (2003).
 - [21] V. Avdeichikov, *et al.*, Nucl. Phys. **A 736**, 22 (2004).
 - [22] C.P. Montoya, *et al.*, Phys. Rev. Lett. **73**, 3070 (1994).
 - [23] J. Töke, *et al.*, Phys. Rev. Lett. **75**, 2920 (1995).
 - [24] Y. Larochelle, *et al.*, Phys. Rev. C **55**, 1869 (1997).
 - [25] J. Lukasik, *et al.*, Phys. Rev. C **55**, 1906 (1997).
 - [26] P. Pawloski, *et al.*, Phys. Rev. C **57**, 1771 (1998).
 - [27] Y. Larochelle, *et al.*, Phys. Rev. C **59**, R565 (1999).
 - [28] E. Plagnol, *et al.*, Phys. Rev. C **61**, 014606 (1999).
 - [29] P.M. Milazzo, *et al.*, Nucl. Phys. **A 703**, 466 (2002).
 - [30] J. Lukasik, *et al.*, Phys. Lett. B, **566**, 76 (2003).
 - [31] G. Lanzanò, *et al.*, Nucl. Phys. **A 683**, 566 (2001).
 - [32] T. Lefort, *et al.*, Nucl. Phys. **A 662**, 397 (2000).
 - [33] W.J. Llope, *et al.*, Phys. Rev. C **52**, 2004 (1995).
 - [34] R. Ghetti, *et al.*, Nucl. Phys. **A674**, 277 (2000).
 - [35] D. R. Tilley, *et al.*, Nucl. Phys. **A 708**, 3 (2002).
 - [36] D. R. Tilley, *et al.*, Nucl. Phys. **A 541**, 1 (1992).
 - [37] R. Lednický, *et al.*, Phys. Lett. B **373**, 30 (1996).
 - [38] R. Ghetti, *et al.*, Phys. Rev. Lett. **87**, 102701-1 (2001).
 - [39] D. Gourio, *et al.*, Eur. Phys. J. A **7**, 245 (2000).

- [40] J. Pochodzalla, *et al.*, Phys. Rev. C **35**, 1695 (1987).
- [41] P. A. DeYoung, *et al.*, Phys. Rev. C **39** (1989) 128; Phys. Rev. C **41** (1990) R1885, and references therein.
- [42] J. Helgesson, *et al.*, submitted to Phys. Rev. C.